


All you need
to know about

DIVERTICULAR DISEASE



Our Digestive System

The Digestive System runs from the mouth to the anus and includes the stomach, the small and large bowels (intestines) and a number of accessory organs. The role of the digestive system is to turn food and liquid into the building blocks that the body needs to function effectively.



Copyright © 2025 This leaflet was published by Guts UK charity in December 2025 and will be reviewed in December 2027. The leaflet was written by Guts UK and reviewed by experts in diverticular disease and has been subject to both lay and professional review. All content in this leaflet is for information only. The information in this leaflet is not a substitute for professional medical care by a qualified doctor or other healthcare professional. We currently use AI translation tools on our website, which may not always provide perfect translations. Please check for further explanation with your doctor if the information is unclear. ALWAYS check with your doctor if you have any concerns about your health, medical condition or treatment. The publishers are not responsible or liable, directly or indirectly, for any form of damages whatsoever resulting from the use (or misuse) of information contained or implied in this leaflet. Please contact Guts UK if you believe any information in this leaflet is in error.




This booklet is about Diverticular Disease of the large bowel (colon)

Diverticular disease is a common condition. It affects around 1 in 3 people, and this increases to 1 in 2 people as they get older. It affects both men and women.

It happens when small pouches form in the wall of the large bowel. These pouches are called diverticula.

What does diverticular disease look like?



Diverticulum: A diverticulum is a small pouch that sticks out from the wall of your large bowel.

Diverticula: This means more than one **Diverticulum**. They are most often found in the lower left part of the bowel.

Diverticulosis: This means you have diverticula in your bowel. Most people with diverticulosis don't have any symptoms and won't need treatment. Many people don't know they have it until they have a test for something else.

Diverticular disease: This means you have diverticula. You might also feel belly pain or notice changes in your bowel habits, like constipation or diarrhoea.

Diverticulitis: This happens when one or more diverticular pouches get inflamed or infected. This can cause increasing pain and other symptoms.

What has caused the development of diverticula?

Diverticula form when the wall of the bowel becomes weak in places, which can happen as we get older. Pressure inside the bowel can then push the lining outwards, forming pouches. Doctors still aren't exactly sure why diverticula form in the bowel. But we do know some things that make it more likely.

Age is the biggest risk factor

Diverticular are more common as you get older and your chances of developing them increases after the age of 40. The number of people affected by diverticula is increasing to 1 in 2 as the population gets older. More people under 45 are now being diagnosed, but this is uncommon.

Other things that may increase the risk

Some lifestyle factors can raise the risk of symptoms or complications. These include:

- Having a higher than healthy body weight.
- Not being physically active.
- Smoking, which is linked to a higher risk of needing hospital care.

What about fibre and constipation?

Some studies suggest that eating more fibre, especially from vegetables, may lower the risk of diverticula. However, other studies find no clear link. It is also **not proven** that constipation causes diverticula.

Genetic conditions (rare causes)

Some rare family conditions can weaken the bowel wall. This makes diverticula more likely.

These include:

- Ehlers-Danlos syndrome
- Williams syndrome
- Polycystic kidney disease
- Marfan syndrome
- Coffin-Lowry syndrome

These conditions can affect the tissue in the bowel, making it weaker and more likely to develop diverticula.



How is diverticular disease diagnosed?

Doctors diagnose diverticular disease by looking inside your bowel or by doing a scan. These tests also help rule out other conditions that could be causing your symptoms.

Tests you may have include:

Colonoscopy or sigmoidoscopy

A thin, flexible tube with a small camera on the end is passed into your bowel through your bottom. This allows the doctor or nurse to see the inside of your large bowel.

A **colonoscopy** looks at the whole bowel, while a **sigmoidoscopy** looks at just the left side of the bowel. It is important that the bowel is empty so that they can see the bowel wall. You will be given a medicine to take at home to make you poo. This medicine will give you loose watery diarrhoea, so you will need to make sure you stay close to a toilet.

CT scan (plain CT)

This is a type of scan that takes pictures of the inside of your belly. It can show signs of diverticular disease and can also help find complications, such as an **abscess** (a pocket of infection).

CT colonography

This is a type of CT scan that looks more closely at the bowel. It's a non-invasive test and doesn't involve putting a camera inside the bowel. A thin tube may be inserted into the rectum to inflate the colon with air, making it easier to see.

Like a colonoscopy, you'll need to **empty your bowels** before the scan using a medicine called a laxative.



How can diverticular disease affect you?

Even with research on this condition, it's still unclear why some people develop complications like infections. Many people can control their symptoms by changing their diet and lifestyle.

Symptoms

Most people with diverticular disease **do not have any symptoms**. Even fewer people will go on to have complications.

Common symptoms

When symptoms do happen, they may include:

- Pain or cramping in your lower belly (usually on the left side).
- A feeling of bloating or swelling in your belly (from gas).
- Changes in your bowel habits. These can include **diarrhoea, constipation**, or switching between both.
- Mucus or blood in your poo.

Everyone's symptoms can be a bit different. The pain may come and go and is often worse after eating. It may feel better after you've been to the toilet for a poo or passed wind. In people of **South Asian background**, the pain is sometimes on the **right side** instead of the left.

Important: Don't ignore new symptoms.

The symptoms of diverticular disease can look like those of other conditions, such as:

- **Irritable bowel syndrome (IBS)**
- **Ulcerative Colitis**
- **Bowel cancer**

So, it's important not to assume that all digestive problems are caused by diverticular disease.

See your GP if you:

- Notice any **new or different symptoms**.
- Have **blood in your poo**.
- Are being sick (vomiting), feel very bloated, have cramping pain and can't poo or pass wind. This may be a **sign of a blockage** or complication, and you should get **urgent medical help**.

Complications

Very few people have complications severe enough to require hospital admission. Also, very few people die from this common disease. However, complications can occur in some cases. These include:

Inflammation (Diverticulitis):

This is the most common complication. The current theory is that the inflammation and/or infection occurs because of an overgrowth of bacteria in the diverticula. It occurs in up to a quarter of people with the condition. In people over 50, acute diverticulitis is more common in females. In those under 50 years of age, it occurs more often in males. More people are being diagnosed, especially people under 45.

Symptoms of diverticulitis:

- Worsening belly pain.
- High temperature.
- Feeling sick and being sick

If these symptoms occur, you should seek medical help immediately.

Complicated diverticulitis requiring hospitalisation is normally treated with antibiotics (germ killing medicine) and fluids. In severe cases, infection can cause complications, such as:

- A collection of pus (abscess).
- An obstruction (blockage of the bowel).
- A perforation (developing a hole in the bowel).

If these complications arise, an operation would be needed to remove part of the large bowel. Or, a needle may be needed to drain a collection of pus.

Blockage of the gut:

Thickening (fibrosis) can occur around areas of diverticulum that have experienced multiple episodes of inflammation. This may lead to narrowing of the large bowel, which can cause a blockage. These narrowings are also referred to as strictures. If diagnosed, surgery is usually needed.

Perforation:

Sometimes, an inflamed pouch (diverticulum) can burst. It causes a hole in the bowel wall called a perforation. A perforation can cause an abscess or more rarely peritonitis. Peritonitis is a serious condition with widespread abdominal inflammation an infection that nearly always requires emergency surgery to treat.

Bleeding:

A blood vessel in a diverticulum can burst. This can cause bleeding in the large bowel. 1 in 3 people with bleeding can have a severe episode. It may require a hospital admission.

Fistulas:

Rarely, abnormal connections can form between diverticula and other organs. Organs affected include the bladder or vagina. These connections are called fistulas.

Symptoms include:

- Poo in urine.
- Foul-smelling vaginal discharge.

If fistulas are diagnosed, surgery is usually necessary. See a doctor if you have these symptoms.

Is there a risk of cancer?

There is **no evidence** that having diverticular disease – whether you have symptoms or not – **increases your risk of bowel cancer**. However, it's still important to be aware of any **new or changing symptoms**. That's because bowel cancer is common in the general population, especially as people get older. Some symptoms of diverticular disease can be like those of bowel cancer.

Always speak to your GP if your symptoms change, get worse, or you notice something new. Especially **blood in your poo, unexplained weight loss, or ongoing changes in your bowel habits**. These symptoms may not be caused by cancer, but it's important to get them checked early.

What treatment is available for diverticular disease?

Even with research on this condition, it's still unclear why some people develop complications like infections. Many people can control their symptoms by changing their diet and lifestyle. The exact cause of flare-ups isn't fully clear. However, a healthy lifestyle may lower the risk of complications.

Diet and lifestyle

Eating well is one of the most important ways to manage diverticular disease.

What to eat

- Follow a **healthy, balanced diet**.
- Include **wholegrains, fruits and vegetables** every day.
- Try to have a **fibre-rich starchy food** (like wholemeal bread, wholegrain pasta, brown rice or potatoes with skin) with each meal.
- Aim for at least **5 portions of fruit and vegetables** per day.
1 portion = about 80g or a handful.
- Smoothies (150ml) count as only **one portion per day**, no matter how much you drink, because they contain less fibre.

Foods containing fibre

This is a list of foods that contain fibre. There are other aspects of healthy eating to consider. Please contact Guts UK for a healthy eating leaflet:
info@gutscharity.org.uk or 020 7486 0341

Starchy foods	Wholemeal, brown or granary bread. Oat or rye bread. Wholegrain crackers, wholegrain rice and wholegrain pasta. Wholegrain breakfast cereals, wheat biscuits and non wholewheat options such as oats, muesli and gluten free wholegrain cereals. Chapatti or naan bread made with wholegrain flour.
Fruit	All varieties are suitable including fresh, frozen, dried and canned.
Vegetables	All varieties are suitable including fresh, frozen, dried and canned. Vegetable soup.
Protein containing foods	Chickpeas, lentils, peas and beans, dhal. Nuts and seeds.
Desserts and other snacks	Puddings containing fruit and dried fruit, oat-based crumble toppings, biscuits containing wholegrain flour. Nuts, dried fruit and seeds, flapjack, muesli bars and fruit bars based on dates. Chocolate containing nuts and dried fruit, root vegetable crisps.
Drinks	Smoothies and milk shakes containing the whole fruit. (150ml portion size. One portion per day counts as one of your 5 fruit and/or vegetables per day)

Drink enough fluids

Try to drink **at least 2 litres** (about 8 to 10 cups) of water or other non-caffeinated drinks each day. This helps fibre move through your bowel and can prevent constipation.

If you're not used to eating much fibre:

Increase fibre **slowly** to avoid bloating or wind. It might take a **few weeks** for your bowel habits to improve.

Do I need to avoid certain foods?

You **don't need to avoid** seeds, nuts, popcorn, or fruit skins. These foods **do not increase the risk** of complications, even if you have diverticular disease.

If you also have irritable bowel syndrome (IBS), eating more wheat bran can make your symptoms worse. If your symptoms don't improve, try increasing your bran fibre intake using non-wheat bran options, like oat or rice bran. These are usually easier for people with IBS to digest.

If you change what you eat, it's important to make sure your diet stays balanced and provides all the nutrients your body needs.

Try to limit red and processed meat to no more than 70 grams per day. If you eat red meat, unprocessed options are a healthier choice.

Red meat: includes beef, lamb, pork, veal, venison, goat, and mutton.

Processed meat: includes sausages, bacon, ham, salami, corned beef, tinned meat, sliced luncheon meats, and pâté.

For more information about healthy eating, fibre, or IBS, contact Guts UK:

Email: info@gutscharity.org.uk

Call: 020 7486 0341

When to ask for more help

You can ask your GP or consultant to refer you to an NHS dietitian if:

- You have another medical condition that affects your diet.
- You're unsure about what foods you should be eating.
- You still have symptoms such as gas, bloating or diarrhoea.

See your GP if your symptoms change.

Other ways to look after your health

- You should take regular exercise [nhs.uk/live-well/exercise/](https://www.nhs.uk/live-well/exercise/)
- If you're overweight, try to lose some weight.
- Stop smoking – it's one of the best things you can do for your health. Stopping smoking can be hard, but getting support makes it easier. Your GP can refer you to a local stop smoking service.

What about probiotics?

Some people try probiotics (good bacteria) to help with diverticular disease. But at the moment, there's not enough good evidence to say they work. So, we don't recommend using probiotics for this condition now.



Medicines

If you have constipation or diarrhoea, speak to your pharmacist or GP. A bulk-forming laxative might help. These work by absorbing water in your bowel, which can help manage both constipation and diarrhoea. If you have pain caused by diverticular disease, your doctor might suggest trying an antispasmodic medicine such as **Mebeverine hydrochloride**.

Avoid using painkillers like **codeine** or other **opioid-based medicines**, as they can make constipation and pain worse. You can read more about this here: gutscharity.org.uk/info/painkillers.

Other medicines that may cause problems include: **Non-steroidal anti-inflammatory drugs (NSAIDs)**, such as ibuprofen and **Nicorandil**, a medicine sometimes used for angina.

Both **opioids** and **NSAIDs** have been linked to serious complications such as **fistulas** and **bowel perforation**.

Important: Do **not** stop taking any prescribed medicines without talking to your doctor. They may be able to offer a safer alternative.

If you have diverticulitis (inflammation)

Your doctor may suggest a 'watch and wait' approach, especially if symptoms are mild. In some cases, antibiotics may be used – for example, if you are unwell, have a weakened immune system, or have other health conditions.

Contact your GP straight away if your symptoms get worse. See the section on **diverticulitis (inflammation)** for more about symptoms.

When is surgery needed?

Surgery for diverticular disease is not usually needed. But it may be offered after a serious episode of inflammation – for example, if:

- An abscess (a collection of pus) forms around the bowel.
- An infection spreads into the abdomen from a perforated (torn) pouch.

In most cases, the operation involves removing the part of the bowel that has the infected or damaged pouches.

Sometimes, the surgeon will need to bring the healthy end of the bowel out through the skin as a **stoma**. This allows waste to leave your body into a bag attached to your tummy. In many cases, this stoma is **temporary** and can be reversed later. This means the bowel can be rejoined if the inflammation has settled and the area has healed well.

Bleeding

Sometimes, diverticular disease can cause bleeding from the bowel. In rare cases, this bleeding can be severe. If this happens, you may need an **emergency blood transfusion**. In most people (8 out of 10), the bleeding **stops on its own** without any treatment.

If the bleeding doesn't stop, your doctor may arrange a **special type of CT scan** to find out where the bleeding is coming from. In many cases, this can also be used to stop the bleeding.

Other treatments using an **endoscope** (a thin, flexible tube with a camera) may also help. Your doctor or nurse will explain which treatment is best for you.

Myths about diverticular disease

You might come across outdated or incorrect advice online about diverticular disease and what foods to avoid. In the past, people were told not to eat certain high-fibre foods like:

- Seeds
- Nuts
- Popcorn
- Fruit skins

It was once believed these foods could get stuck in the diverticula (small pouches in the bowel) and cause problems. **But this is no longer thought to be true.** These foods **do not** increase the risk of diverticulitis or bleeding. In fact, research shows:

- **Men who eat nuts and popcorn** had a **lower risk** of diverticulitis and bleeding.
- **Women who eat more fibre** were **less likely** to develop diverticulitis.

So, unless your doctor or dietitian tells you otherwise, these foods can be enjoyed as part of a healthy, balanced diet.

Myths about diverticulitis

Some people are told to eat less fibre during a flare-up of **diverticulitis** (inflammation or infection of the diverticula). But recent reviews show that there's **no good evidence** that a **low-fibre or low-residue** diet helps people recover faster, have fewer symptoms, or avoid future episodes. **A normal, balanced diet** may be just as effective for people with **uncomplicated diverticulitis**.

If your diverticulitis becomes more serious – for example, if you develop **a fistula, abscess, bowel blockage, perforation**, or if you've had **surgery** – your doctor or dietitian may give you different advice based on your specific needs.

What to ask your doctor

It's ok to have questions about your condition and treatment. These are some useful questions you can ask your doctor:

- *Has my colonoscopy/CT scan ruled out any other conditions?*
- *Are there any issues of potential concern in my large bowel?*
- *Would you refer me to a dietitian for further help with my diet?*
- *What monitoring or follow up will I be offered?*
- *When will antibiotics be considered when treating my diverticulitis and when should I seek further advice?*



Where can I get more information about treatment?

You can find reliable, evidence-based information from the **National Institute for Health and Care Excellence (NICE)**. They provide clear, evidence-based guidance on a wide range of health, public health, and social care topics. NICE provides guidance on:

- How to prevent and manage different health conditions.
- The most effective treatments and medicines.
- Ways to improve health and care in various settings.

Their recommendations are based on the best available research and are used by the NHS to help guide treatment and care.

You can visit the NICE website to learn more: [nice.org.uk/guidance/ng147](https://www.nice.org.uk/guidance/ng147)

References

References are available on request.

Contact us on **0207 486 0341** or email info@gutscharity.org.uk for more information.

How can I be involved in research into diverticular disease?

Guts UK funded a Priority Setting Partnership (PSP) into diverticular disease to help identify the most important areas for research into the condition. Healthcare professionals, patients and their families identified the top ten priorities for future research. Find out more at gutscharity.org.uk/psps.

It is important that research into the causes and treatments for diverticular disease continues. Guts UK is the only UK charity funding research into the whole digestive system. We're proud to provide patient information about diverticular disease and are here to help.

Please contact gutscharity.org.uk/contact-us/ with any questions that you may have.

If you found this information useful, please consider donating to support our work into diverticular disease today. Thank you.
gutscharity.org.uk/donate/

Guts UK

is the national charity for the digestive system



Our three main areas of work are:

- Information and support
- Raising awareness and public education
- Research with patient and public involvement and engagement (PPIE)

Guts UK provides information and support to people affected by digestive conditions and symptoms. We raise vital awareness about our guts and fund life-changing research into the digestive system. Our mission is a world where digestive conditions are better understood, better treated and everyone who lives with one gets the support they need.

Get in touch

If you need information and support for digestive conditions or symptoms, please call our freephone Helpline on **0300 102 4887** (Monday to Friday, 10am to 2pm).

Alternatively you can use the online form on our website at www.gutscharity.org.uk/helpline.

For general enquiries about anything else, please call us on **0207 486 0341** or email info@gutscharity.org.uk. You can find out more about Guts UK at www.gutscharity.org.uk.



 **@GutsCharityUK**

At Guts UK we only want to send you information you want to receive, the way you want to receive it. We take great care of your personal data and never sell or swap data. Our privacy policy is online at www.gutscharity.org.uk and you can always change your preferences by contacting us at info@gutscharity.org.uk or calling 0207 486 0341.

Linda's diverticular disease story

"I didn't want the pain to control my life anymore."

For years, Linda lived with the unpredictable and debilitating pain of diverticular disease. What started as occasional discomfort soon turned into severe flare-ups that left her feeling exhausted and isolated. Simple joys, like going out for a meal or traveling, became sources of intense anxiety.

After a frightening emergency admission to the hospital, Linda knew she needed to find a better way to manage her condition. With the right information and support, she began to understand her triggers and take back control of her health.

But no one should have to suffer in silence. Research into diverticular disease is vital to understanding why flare-ups happen and how we can prevent them. By better understanding the gut, we can ensure that people like Linda can live their lives to the fullest, free from the fear of the next attack.

When you support Guts UK, you fund life-changing research and information that provides hope and clarity for the millions of people living with digestive conditions.



Donation Form

I would like to make a donation to Guts UK.



Title First name
Surname
Address
 Postcode
Tel Email
Guts UK Reference: 0001

If you wish, please share with us your motivation for giving today. This will help us tailor our thank you:

Please tick here if you do not wish to receive a thank you letter to acknowledge your donation.

I would like to support Guts UK with a donation of

£5 £10 £25 £50 £100 £500 Other £

I enclose a cheque payable to Guts UK OR Please debit my credit/debit card

Card no. Expiry CVC / Security code

Address
(if different
from above)

OR to take my details

Signature(s) Date

We are incredibly grateful for all donations made to Guts UK in support of our work. We will write to thank you for your donation but understand that sometimes, donors prefer not to receive this kind of communication. Please tick here if you do not wish to receive a thank you.

Please turn every £10 I donate into £12.50 at no extra cost to me, by adding gift aid to my donation. Add Gift Aid
I am a UK taxpayer, please treat all donation I make or have made to Guts UK in the past 4 years as Gift Aid donations until further notice.

For more information on Gift Aid please see below.

Signature(s) Date

giftaid it I am happy for all gifts of money that I have made to Guts UK charity (Core) in the last four years and all future gifts of money that I make to be Gift Aid donations. I am a UK taxpayer and understand that if I pay less Tax & Capital Gains Tax in that year that the amount of Gift Aid claimed on all my donations across all charities, it is my responsibility to pay any difference. Guts UK charity claims 25p for every £1 you donate from the tax you pay for the current tax year. If your circumstances, name or address change please do let us know.

Welcome to Guts UK

Information is power. Armed with information, patients can make informed decisions and take control. Choose how you can stay in touch with Guts UK and keep up to date with our latest information and research:

By email
 By post
 By telephone

Please return this donation form to this FREEPOST address:
Freepost GUTS-UK-CHARITY

Alternatively, you can:
Call us on **020 7486 0341**
or go to gutscharity.org.uk/donate.